

Long Waves Dissipation and Harmonic Generation by Coastal Forests

- Che-Wei Chang¹, Philip L.-F. Liu^{2,3}
- August 1st, 2018
- School of Civil and Environmental Engineering
- Cornell University

¹ Now at: Disaster Prevention Research Institute, Kyoto University

² Department of Civil and Environmental Engineering, National University of Singapore

³ Institute of Hydrological and Oceanic Sciences, National Central University



Motivations & Objectives

- Coastal forests – **non-intrusive (natural) protection** against ocean waves
- How effective can coastal vegetation dissipate incoming wave energy?
- Interactions between waves and vegetation:
 - Physical modeling – rigid/flexible cylinders or live vegetation (Wu et al. 2011, Maza et al. 2015)
 - Numerical modeling – N-S models, depth-integrated models (NLSW, Boussinesq-type equations)
 - Mathematical modeling – Homogenization theory (Mei et al. 2011, 2014)
- Previous work:
 - Develop a model to estimate wave attenuation by coastal forests of arbitrary shape
 - Linear model vs. experimental data (Liu et al. 2015, Chang et al. 2017a, b)



How about nonlinearity?




Motivations & Objectives


- Previous work:
 - Develop a model to estimate wave attenuation by coastal forests of arbitrary shape
 - Linear model vs. experimental data (Liu et al. 2015, Chang et al. 2017a, b)

How about nonlinearity?

-
- Extend the linear model (Mei et al. 2011) – homogenization theory
 - Consider the effects of weak nonlinearity
 - Investigate the nonlinear effects and harmonic generation



Periodic shallow-water waves




- Conditions:

- Shallow water: wavelength is much greater than water depth
- Tree spacing is much smaller than the wavelength
- Incident waves: simple-harmonic waves with weak nonlinearity

➔ $\ell \ll h_0 \ll 1/k_{inc}$



Periodic shallow-water waves



- Governing equations (shallow water):

$$\frac{\partial \eta}{\partial t} + \frac{\partial}{\partial x_i} [u_i (h_0 + \eta)] = 0, \quad i = 1, 2$$

$$\frac{\partial u_i}{\partial t} + \boxed{u_j \frac{\partial u_i}{\partial x_j}} = -g \frac{\partial \eta}{\partial x_i} + \nu_e \frac{\partial^2 u_i}{\partial x_j \partial x_j}, \quad i \& j = 1, 2$$

u_i : velocity components

η : free surface elevation


ν_e : eddy viscosity ← Spatial average

Incident waves:

simple-harmonic waves with weak nonlinearity



Periodic shallow-water waves



- Governing equations (shallow water):

$$\frac{\partial \eta}{\partial t} + \frac{\partial}{\partial x_i} [u_i (h_0 + \eta)] = 0, \quad i = 1, 2$$

$$\frac{\partial u_i}{\partial t} + \boxed{u_j \frac{\partial u_i}{\partial x_j}} = -g \frac{\partial \eta}{\partial x_i} + \nu_e \frac{\partial^2 u_i}{\partial x_j \partial x_j}, \quad i \& j = 1, 2$$

u_i : velocity components

η : free surface elevation

ν_e : eddy viscosity ← Spatial average


- Parameters:

$$\varepsilon = k_{inc} \ell = \frac{\omega \ell}{\sqrt{g h_0}} \ll \mathcal{O}(1), \quad \alpha = \frac{H_{inc}/2h_0}{\varepsilon} = \mathcal{O}(1) \Rightarrow \boxed{\mathcal{O}\left(\frac{H_{inc}}{2h_0}\right) = \mathcal{O}(\varepsilon)}$$

Weakly nonlinear waves



Periodic shallow-water waves



- Governing equations (shallow water):

$$\frac{\partial \eta}{\partial t} + \frac{\partial}{\partial x_i} [u_i (h_0 + \eta)] = 0, \quad i = 1, 2$$

$$\frac{\partial u_i}{\partial t} + \boxed{u_j \frac{\partial u_i}{\partial x_j}} = -g \frac{\partial \eta}{\partial x_i} + \nu_e \frac{\partial^2 u_i}{\partial x_j \partial x_j}, \quad i \& j = 1, 2$$

u_i : velocity components

η : free surface elevation

ν_e : eddy viscosity ← Spatial average

- Eddy viscosity (Mei et al. 2011):


$$\nu_e = 1.86(1 - n)^{2.06} U_0 \ell, \quad U_0 = \sqrt{gh_0} A_{inc} / h_0$$

porosity

Shallow-water wave characteristic velocity



Homogenization (multi-scale perturbation theory)



Macro-scale Problem
 < scale: wavelength >

$$X_i^* = \varepsilon x_i^*$$

Micro-scale Problem
 < scale: tree spacing >

$$x_i^* = x_i / \ell$$

dimensionless

$$\varepsilon = k_{inc} \ell = \frac{\omega \ell}{\sqrt{g h_0}} \ll \mathcal{O}(1)$$

$$(u_i^*, \eta^*) = (u_i^{*(0)}, \eta^{*(0)}) + \varepsilon (u_i^{*(1)}, \eta^{*(1)}) + \dots \implies u_i^{*(n)}, \eta^{*(n)} : \text{functions of } (x_i^*, X_i^*, t^*)$$



Leading-order problem

- o Summation of different harmonics

$$u_i = \frac{1}{2} \sum_{m=-\infty}^{\infty} \tilde{u}_{i,m} e^{-imt} \quad \eta = \frac{1}{2} \sum_{m=-\infty}^{\infty} \tilde{\eta}_m e^{-imt} \quad \longrightarrow \quad (\tilde{u}_{i,-m}, \tilde{\eta}_{-m}) = \text{complex conjugate of } (\tilde{u}_{i,m}, \tilde{\eta}_m)$$

- Micro-scale (cell) problem - **NONLINEAR**

$$\frac{\partial \tilde{u}_{i,m}^{(0)}}{\partial x_i} = 0, \quad \vec{x} \in \Omega_f$$


$$(-im) \tilde{u}_{i,m}^{(0)} + \frac{\alpha}{2} \sum_{m_1=-\infty}^{\infty} \left(\tilde{u}_{j,m_1}^{(0)} \frac{\partial \tilde{u}_{i,m-m_1}^{(0)}}{\partial x_j} \right) = -\frac{\partial \tilde{\eta}_m^{(0)}}{\partial X_i} - \frac{\partial \tilde{\eta}_m^{(1)}}{\partial x_i} + \sigma \frac{\partial^2 \tilde{u}_{i,m}^{(0)}}{\partial x_j \partial x_j}, \quad \vec{x} \in \Omega_f$$

Harmonic generation

Macro-scale pressure gradient

Dimensionless eddy viscosity:

$$\sigma = \frac{\nu_e}{\omega \ell^2} \equiv 1.86(1-n)^{2.06} \frac{1}{k_{inc} \ell} \left(\frac{A_{inc}}{h_0} \right), \quad k_{inc} = \frac{\omega}{\sqrt{gh_0}}$$



- Boundary conditions:

$$\tilde{u}_{i,m}^{(0)} = 0, \quad \vec{x} \in S_c$$

$$\langle \tilde{\eta}_m^{(1)} \rangle = \frac{1}{\Omega} \iint_{\Omega_f} \tilde{\eta}_m^{(1)} dx_1 dx_2 = 0$$

Nonlinear B.V.P – Unknowns: $\tilde{u}_{i,m}^{(0)}$ and $\tilde{\eta}_m^{(1)}$



Leading-order problem

- Micro-scale (cell) problem - **NONLINEAR**


➤ Modified pressure correction method – iteration

Dimensionless eddy viscosity

$$\frac{\partial \tilde{u}_{i,m}^{(0)}}{\partial t} - \text{im} \tilde{u}_{i,m}^{(0)} + \frac{\alpha}{2} \sum_{m_1=-\infty}^{\infty} \frac{\partial \left(\tilde{u}_{j,m_1}^{(0)} \tilde{u}_{i,m-m_1}^{(0)} \right)}{\partial x_j} = - \frac{\partial \tilde{\eta}_m^{(0)}}{\partial X_i} - \frac{\partial \tilde{\eta}_m^{(1)}}{\partial x_i} + \sigma \frac{\partial^2 \tilde{u}_{i,m}^{(0)}}{\partial x_j \partial x_j}$$

Pseudo-time derivative

Macro-scale pressure gradient (GIVEN)



➤ Finite difference with staggered discretization:

$$\frac{\left(\tilde{u}_{i,m}^{(0)} \right)^{n_t+1} - \left(\tilde{u}_{i,m}^{(0)} \right)^{n_t}}{\Delta t} = \text{im} \left(\tilde{u}_{i,m}^{(0)} \right)^{n_t} - \frac{\alpha}{2} \sum_{m_1=-\infty}^{\infty} \left(\frac{\partial \tilde{u}_{j,m_1}^{(0)} \tilde{u}_{i,m-m_1}^{(0)}}{\partial x_j} \right)^{n_t} - \frac{\partial \tilde{\eta}_m^{(0)}}{\partial X_i} - \left(\frac{\partial \tilde{\eta}_m^{(1)}}{\partial x_i} \right)^{n_t} + \sigma \left(\frac{\partial^2 \tilde{u}_{i,m}^{(0)}}{\partial x_j \partial x_j} \right)^{n_t}$$

if convergent

→ $\left(\tilde{u}_{i,m}^{(0)} \right)^{n_t+1} \approx \left(\tilde{u}_{i,m}^{(0)} \right)^{n_t}$ $(n_t)^{\text{th}}$ iteration



Leading-order problem

- Macro-scale (wavelength-scale) problem

- Forest region for each harmonic – **LINEAR**

$$n \left(-im\tilde{\eta}_m^{(0)} \right) + \frac{\partial \langle \tilde{u}_{i,m}^{(0)} \rangle}{\partial X_i} = 0 \quad \leftarrow \text{cell-averaged quantity}$$

$$-im \langle \tilde{u}_{i,m}^{(0)} \rangle + \alpha \tilde{M}_m = -n \frac{\partial \tilde{\eta}_m^{(0)}}{\partial X_i} - \tilde{N}_m + \sigma \tilde{Q}_m$$

$$\Rightarrow \frac{\partial^2 \tilde{\eta}_m^{(0)}}{\partial X_i^2} + (m^2) \tilde{\eta}_m^{(0)} = - \left(\frac{\alpha}{n} \right) \frac{\partial \tilde{M}_m}{\partial X_i} - \left(\frac{1}{n} \right) \frac{\partial \tilde{N}_m}{\partial X_i} + \left(\frac{\sigma}{n} \right) \frac{\partial \tilde{Q}_m}{\partial X_i}$$

- Complex coefficients:

Cell problem solutions

$$\tilde{M}_m = \frac{1}{2\Omega} \iint_{\Omega_f} \left[\sum_{m_1=-\infty}^{\infty} \tilde{u}_{j,m_1}^{(0)} \frac{\partial \tilde{u}_{i,m-m_1}^{(0)}}{\partial x_j} \right] d\Omega, \quad \tilde{N}_m = \frac{1}{\Omega} \iint_{\Omega_f} \frac{\partial \tilde{\eta}_m^{(1)}}{\partial x_i} d\Omega, \quad \tilde{Q}_m = \frac{1}{\Omega} \iint_{\Omega_f} \frac{\partial^2 \tilde{u}_{i,m}^{(0)}}{\partial x_j \partial x_j} d\Omega$$

- Open water region for each harmonic – **LINEAR**

$$-im\tilde{\eta}_m^{(0)} + \frac{\partial \langle \tilde{u}_{i,m}^{(0)} \rangle}{\partial X_i} = 0$$

$$-im \langle \tilde{u}_{i,m}^{(0)} \rangle = - \frac{\partial \tilde{\eta}_m^{(0)}}{\partial X_i}$$

$$\Rightarrow \frac{\partial^2 \tilde{\eta}_m^{(0)}}{\partial X_i^2} + (m^2) \tilde{\eta}_m^{(0)} = 0$$



Leading-order problem


- Micro-scale problem: $\frac{\partial \tilde{u}_{i,m}^{(0)}}{\partial x_i} = 0, \quad \vec{x} \in \Omega_f$

$$(-im) \tilde{u}_{i,m}^{(0)} + \frac{\alpha}{2} \sum_{m_1=-\infty}^{\infty} \left(\tilde{u}_{j,m_1}^{(0)} \frac{\partial \tilde{u}_{i,m-m_1}^{(0)}}{\partial x_j} \right) = - \frac{\partial \tilde{\eta}_m^{(0)}}{\partial X_i} - \frac{\partial \tilde{\eta}_m^{(1)}}{\partial x_i} + \sigma \frac{\partial^2 \tilde{u}_{i,m}^{(0)}}{\partial x_j \partial x_j}, \quad \vec{x} \in \Omega_f$$

Macro-scale pressure gradient

- Macro-scale problem: $\frac{\partial^2 \tilde{\eta}_m^{(0)}}{\partial X_i^2} + (m^2) \tilde{\eta}_m^{(0)} = - \left(\frac{\alpha}{n} \right) \frac{\partial \tilde{M}_m}{\partial X_i} - \left(\frac{1}{n} \right) \frac{\partial \tilde{N}_m}{\partial X_i} + \left(\frac{\sigma}{n} \right) \frac{\partial \tilde{Q}_m}{\partial X_i}$

Cell problem solutions



Long waves through a forest belt

- Macro-scale (wavelength-scale) problem

➤ Forest region:
$$\frac{\partial^2 \tilde{\eta}_m^{(0)}}{\partial X^2} + (m^2) \tilde{\eta}_m^{(0)} = - \left(\frac{\alpha}{n} \right) \frac{\partial \tilde{M}_m}{\partial X} - \left(\frac{1}{n} \right) \frac{\partial \tilde{N}_m}{\partial X} + \left(\frac{\sigma}{n} \right) \frac{\partial \tilde{Q}_m}{\partial X}, \quad \text{if } 0 < X < L_F$$

➤ Open water:
$$\frac{\partial^2 \tilde{\eta}_m^{(0)}}{\partial X^2} + (m^2) \tilde{\eta}_m^{(0)} = 0, \quad \text{if } X < 0 \quad \text{or} \quad X > L_F$$

$$\tilde{\eta}_{I,m}^{(0)} = \mathcal{I}_m e^{imX} + \mathcal{R}_m e^{-imX}, \quad \tilde{u}_{I,m}^{(0)} = \mathcal{I}_m e^{imX} - \mathcal{R}_m e^{-imX} \quad \text{if } X < 0$$

$$\tilde{\eta}_{T,m}^{(0)} = \mathcal{T}_m e^{imX}, \quad \tilde{u}_{T,m}^{(0)} = \mathcal{T}_m e^{imX} \quad \text{if } X > L_F$$


$$\mathcal{R}_0 = 0$$

➔ Mean water level of incidence region = constant

➤ Matching conditions:

$$\eta_{I,m}^{(0)} = \eta_{F,m}^{(0)}, \quad \tilde{u}_{I,m}^{(0)} = \tilde{u}_{F,m}^{(0)} \quad \text{at } X = 0$$





$$\eta_{F,m}^{(0)} = \eta_{T,m}^{(0)}, \quad \tilde{u}_{F,m}^{(0)} = \tilde{u}_{T,m}^{(0)} \quad \text{at } X = L_F$$



Long waves through a forest belt

- **Case 1:** $T = 2.50$ s, $A_{inc} = 0.33$ cm
 $h_0 = 12$ cm, $L = 2.692$ m
 $\alpha = 0.3957$, $\sigma = 0.0048$





Circles — linear model solution
 Solid line — upper wave envelope



Long waves through a forest belt

- **Case 2:** $T = 1.90$ s, $A_{inc} = 1.22$ cm
 $h_0 = 12$ cm, $L = 2.038$ m
 → $\alpha = 1.1119$, $\sigma = 0.0136$



Circles — linear model solution
 Solid line — upper wave envelope



Long waves through a forest belt

$$\alpha = \frac{A_{\text{inc}}/h_0}{k_{\text{inc}}\ell} = \frac{A_{\text{inc}}/h_0}{\varepsilon}$$

$$\sigma = 1.86(1 - n)^{2.06} \frac{1}{k_{\text{inc}}\ell} \left(\frac{A_{\text{inc}}}{h_0} \right) \equiv 1.86(1 - n)^{2.06} \cdot \alpha$$



Reflection coefficient: $\bar{R} = \left(\sum_{m=0}^{\infty} |\mathcal{R}_m|^2 \right)^{1/2}$



Transmission coefficient: $\bar{T} = \left(\sum_{m=0}^{\infty} |\mathcal{T}_m|^2 \right)^{1/2}$



Long waves through a forest belt

$$\alpha = \frac{A_{\text{inc}}/h_0}{k_{\text{inc}}\ell} = \frac{A_{\text{inc}}/h_0}{\varepsilon}$$

$$\sigma = 1.86(1-n)^{2.06} \frac{1}{k_{\text{inc}}\ell} \left(\frac{A_{\text{inc}}}{h_0} \right) \equiv 1.86(1-n)^{2.06} \cdot \alpha$$



Reflection coefficient: $\bar{R} = \left(\sum_{m=0}^{\infty} |\mathcal{R}_m|^2 \right)^{1/2}$

Transmission coefficient: $\bar{T} = \left(\sum_{m=0}^{\infty} |\mathcal{T}_m|^2 \right)^{1/2}$



Summary

- Micro-scale problem:
 - The numerical model for solving micro-scale nonlinear problem is developed
 - Boundary-fitting discretization is needed for improvement
- Macro-scale problem:
 - Higher harmonics are generated and radiated into outside region
 - The first harmonic is dominant and higher harmonics have smaller amplitude
 - Lack of gauge data for shallow-water waves
 - Model extension for taking vertical variation into account is needed
 - The use of drag coefficient and eddy viscosity will be made
 - More gauge data are available for model validation





36TH INTERNATIONAL CONFERENCE ON COASTAL ENGINEERING 2018

Baltimore, Maryland | July 30 – August 3, 2018

The State of the Art and Science of Coastal Engineering

Thank You

Che-Wei Chang

cc2338@cornell.edu

